

less impressed, although more so than in *Wheeleri* Mel.: front tibiae with a suffused broad dark band.

The females of *kansensis* and *trinitatus* and markedly alike, but the males are entirely different. Dr. Adams suggested in a letter the possibility of a species with dimorphic males, but a close study reveals characters by which the females can be recognized, which indicates that there are two distinct species. The differences in the extent of the color markings of the abdomen and legs are not of value in separating these species. The characteristic differences between the two species may be stated thus:

Nemotelus kansensis Adams.

Female: 5.5 mm. Rostellum projecting beyond the eye more than the horizontal diameter of the eye: proboscis geniculate a little before the middle.

Nemotelus trinitatus Melander.

Female: Length 4.5 mm. Rostellum projecting not more than the diameter of the eye: proboscis geniculate at the middle.

The male of *kansensis* is at once distinguished from all the other species by the single conspicuous black fascia on the fifth abdominal segment. *Slossonae* Johnson and *flavicornis* Johnson, the only other species with a black fascia so placed, are of small size and have the fourth segment also blackened. Moreover in these species the third vein is simple.

A DIPTEROUS PARASITE OF THE BOX TURTLE.—In PSYCHE, Vol. V, page 403, Dr. Wm. M. Wheeler mentions several cases of finding larvae of dipterous flies of the genus *Sarcophoga* in tumors in the skin of the Box Turtle. On July 28, 1902, I found another case of the same kind at Cold Spring Harbor, Long Island, N. Y., near the biological laboratory. The turtle had a swelling about an inch in diameter on the left side of the neck with a small opening directed forward that was usually nearly closed but could be easily stretched to quarter of an inch in diameter. Five larvae were taken out through this opening with forceps, one dead and partly decayed, the others alive and full grown. Placed in bottles with moist earth they buried themselves within a few hours. On July 31 one of them had pupated and the fly came out August 17. It is plainly a *Sarcophoga* but has not yet been examined by anyone familiar enough with this genus to determine the species. The fly and one of the larvae are in the Museum of Comparative Zoölogy in Cambridge, Mass.—*J. H. Emerton.*